

ACCEPTED MANUSCRIPT • OPEN ACCESS

Cocoa agroforestry systems versus monocultures: a multi-dimensional meta-analysis

To cite this article before publication: Wiebke Niether *et al* 2020 *Environ. Res. Lett.* in press <https://doi.org/10.1088/1748-9326/abb053>

Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2020 The Author(s). Published by IOP Publishing Ltd.

As the Version of Record of this article is going to be / has been published on a gold open access basis under a CC BY 3.0 licence, this Accepted Manuscript is available for reuse under a CC BY 3.0 licence immediately.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected and is not published on a gold open access basis under a CC BY licence, unless that is specifically stated in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **Cocoa agroforestry systems versus monocultures: a multi-**
2 **dimensional meta-analysis**

3 Wiebke Niether^{1,*}, Johanna Jacobi^{2,*}, Wilma J. Blaser³, Christian Andres³, and Laura Armengot⁴

4 ¹Institute of Geography, University of Göttingen, 37077 Göttingen, Germany

5 ²Centre for Development and Environment, University of Bern, 3012 Bern, Switzerland

6 ³Department of Environmental Systems Science, ETH Zurich, 8092 Zurich, Switzerland

7 ⁴International Cooperation Department, Research Institute of Organic Agriculture, FiBL,
8 Switzerland

9 * Wiebke Niether; Johanna Jacobi.

10 Email: wiebke.niether@geo.uni-goettingen.de; johanna.jacobi@cde.unibe.ch

Abstract

Scientific knowledge, societal debates, and industry commitments around sustainable cocoa are increasing. Cocoa agroforestry systems are supposed to improve the sustainability of cocoa production. However, their combined agronomic, ecological, and socio-economic performance compared to monocultures is still largely unknown. Here we present a meta-analysis of 52 articles that directly compared cocoa agroforestry systems and monocultures. Using an inductive, multi-dimensional approach, we analyzed the differences in cocoa and total system yield, economic performance, soil chemical and physical properties, incidence of pests and diseases, potential for climate change mitigation and adaptation, and biodiversity conservation. Cocoa agroforestry systems outcompeted monocultures in most indicators. Cocoa yields in agroforestry systems were 25% lower than in monocultures, but total system yields were about 10 times higher, contributing to food security and diversified incomes. This finding was supported by a similar profitability of both production systems. Cocoa agroforestry contributed to climate change mitigation by storing 2.5 times more carbon and to adaptation by lowering mean temperatures and buffering temperature extremes. We found no significant differences in relation to the main soil parameters. The effect of the type of production system on disease incidence depended on the fungal species. The few available studies comparing biodiversity showed a higher biodiversity in cocoa agroforestry systems. Increased and specific knowledge on local tree selections and local socio-economic and environmental conditions, as well as building and enabling alternative markets for agroforestry products, could contribute to further adoption and sustainability of cocoa agroforestry systems.

Keywords

Economic performance; system yield; pests and diseases; biodiversity; sustainability; *Theobroma cacao*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1. Introduction

Cocoa is an important commodity worldwide. In 2018, up to 5.3 million metric tons of dry cocoa beans were produced on about 12 million hectares [1], mostly by smallholders [2]. The 2018 market value of processed cocoa was estimated at USD 13.4 billion [3]. The increasing global demand for cocoa has led to intensification of cocoa production systems. Consequently, cultivation of cocoa under tree shade is gradually being replaced with full-sun monocultures and the use of agrochemicals [4-6]. Monocultures are often reported to have higher cocoa yields than agroforestry systems, but they cause adverse social-ecological impacts [7, 8]. In many areas, intensified cocoa production has led to deforestation, biodiversity loss, increased carbon emissions, reduction of energy efficiency, soil degradation, and contamination from pesticides [9, 10] as well as to socio-economic problems such as food insecurity and vulnerability to cocoa price volatilities [11]. Unsustainable exploitation of natural resources can lead to environmental disasters and social conflicts [12], while crop diversification strategies aim to reduce environmental impacts [13]. Therefore, farmers' livelihoods, including both profitability and food security, need to be considered together with other sustainability indicators of cocoa production systems such as biodiversity conservation, climate change mitigation and adaptation, and soil fertility.

Increasing yields while reducing costs is one of the main targets of the cocoa industry. Producers need to make a living from cocoa as a cash crop, and are often circumspect of cocoa agroforestry systems due to the likelihood of increased costs of labor and inputs. Lack of knowledge on the management of shade and fruit trees, access to planting material or tools,

and market constraints are some of the reasons that prevent a broader adoption of agroforestry systems [14]. However, the growing social awareness of cocoa's social-ecological impacts is putting pressure on the cocoa value chain to source from sustainable production systems that minimize deforestation, biodiversity degradation and child labor, and allow farmers to earn a living income. The sustainability commitments of some of the largest chocolate companies indicate the need and willingness of the processing industry to invest in the sustainability of the cocoa value chains [15, 16]. Agroforestry systems, which grow timber, fruit and other trees together with cocoa trees, have the potential to increase the sustainability of cocoa production. Trees on agricultural land play a crucial role in a context of climate change through carbon sequestration in biomass and soils [8, 13], as well as in adapting to climate change [7, 17] by buffering climatic extremes and blocking direct radiation [7, 18]. Agroforestry systems can provide a variety of habitats and microclimates that support biodiversity conservation [19-21]. Both microclimate and biodiversity can regulate the incidence of pests and diseases [22, 23]. Trees may improve the functional diversity, nutrient cycling, and soil chemical and physical properties of cocoa production systems [17]. Furthermore, trees can provide additional economic return, e.g. from fruit or timber [24], and increase local food security [5, 25]. In the long-term, cocoa agroforestry systems may even provide higher cocoa yields than monocultures due to an observed early aging of unshaded cocoa trees [26, 27].

Although many studies describe the benefits of cocoa agroforestry systems, a quantitative consolidation of the benefits and drawbacks of cocoa agroforestry systems in direct comparison with cocoa monocultures is lacking. De Beenhouwer et al. [28] compared studies on cocoa and coffee agroforestry systems with natural forest and plantations with sparse shade trees, but full-sun monocultures were not included. Therefore, our aim was to conduct a meta-analysis on the performance of cocoa agroforestry systems compared to monocultures, including the most

1
2
3 86 studied key indicators, i.e., yield, economic performance, soil fertility, pests and diseases,
4
5 87 carbon sequestration, microclimate, and biodiversity conservation. In particular, we addressed
6
7 88 the questions whether 1) cocoa agroforestry systems increase productivity; 2) cocoa
8
9 89 agroforestry systems sustain farmers' incomes; 3) cocoa agroforestry systems improve soil
10
11 90 chemical and physical properties for cocoa production; 4) cocoa agroforestry systems enhance
12
13 91 the control of pests and diseases; 5) cocoa agroforestry systems support the adaptation of
14
15 92 cocoa plantations to climate change, 6) cocoa agroforestry systems contribute to climate change
16
17 93 mitigation; and 7) cocoa agroforestry systems contribute to biodiversity conservation in
18
19 94 comparison to cocoa monocultures.
20
21
22
23

24 95 **2. Methodology**

25 96
26 97 *2.1 Literature selection*

27
28
29 98 We gathered scientific peer-reviewed articles from Web of Science in February 2020 by
30
31 99 searching with the keyword combinations "(TS = (cacao OR cocoa) AND agroforest*)" and "(TS =
32
33 100 (cacao OR cocoa) AND *shade*)", where TS refers to topics mentioned in the title and abstract
34
35 101 of the articles. In this meta-analysis, we focused on peer-reviewed articles in English. We
36
37 102 discarded articles that did not report information or results related to the production system.
38
39 103 We completed the database with already collected publications about cocoa agroforestry
40
41 104 systems drawn from our own libraries (66 articles). This resulted in a total number of 542
42
43 105 articles on cocoa production systems. Figures S3 and S4 (SI Appendix) provide, respectively, an
44
45 106 overview of the articles by country and year. We screened all articles for their suitability to be
46
47 107 included in the meta-analysis. We excluded 28 articles that were not reporting original data
48
49 108 (meta-analyses and reviews), nine articles that were based on modelling, and two studies on
50
51 109 cocoa grown below a shade roof instead of a natural tree canopy. A large number of articles
52
53
54
55
56
57
58
59
60

(420) were not included because they did not compare cocoa agroforestry systems with cocoa monocultures. 21 articles that compared cocoa agroforestry systems with monocultures were excluded due to a lack of information (on sample size, means or standard deviations) or covered topics that were not included in this meta-analysis. Six more studies did not provide quantitative data, and four studies were not accessible. Finally, we analyzed 52 articles presenting results from cocoa farms or experimental stations (SI Appendix, Table S1). The research presented in these articles covers three continents and ten countries, i.e. Ghana (20), Cameroon (2), Ivory Coast (1), Indonesia (8), Malaysia (2), Costa Rica (1), Panama (1), Ecuador (3), Peru (1) and Bolivia (13). Ivory Coast, the world's leading cocoa producing country, is represented with only one article, due to very few studies published in English, compared to the high amount of research done in Ghana by the Cocobod, national and international institutions (SI Appendix, Figure S4). In contrast, many data come from the rather small cocoa producing country Bolivia due to the existence of a long-term field trial comparing different cocoa production systems. The first article comparing cocoa agroforestry systems with cocoa monocultures was published in 1968, but 70 % of the articles included were published after 2010.

2.2 Limitations of the study

The restriction of our analysis to studies published in English may have led to the exclusion of important research in other languages. However, to the best of our knowledge, a scientific database for non-English publications comparable to Web of Science does not exist at present, impeding a systematic search for peer-reviewed publications. The qualitative studies excluded from our meta-analysis addressed aspects such as livelihoods, cultural and social services, and political issues. These topics are therefore underrepresented in our study. Due to their high

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

relevance for the performance of cocoa production systems, they should be covered in future analyses.

It became evident from our literature search that a clear definition of cocoa agroforestry systems does not exist and that cocoa agroforestry systems span across a wide range of designs and management activities (SI Appendix, Table S2). This variation has to do with differences in ecosystems, planting material (selection of and access to tree species and cocoa varieties), farmers and their culture, local knowledge, market conditions, access to information and tools, soil and climatic conditions, landscapes and land-use histories. The aim of our study was to compare cocoa agroforestry systems with cocoa monocultures, while aware of the heterogeneity of cocoa agroforestry systems; we assumed that even a simple cocoa agroforestry system could affect the analyzed factors and reveal a difference with cocoa monocultures [7]. In addition, cocoa monocultures also vary due to environmental conditions, management practices, and the selection of cocoa varieties. However, not all the articles provided detailed information on these aspects. Therefore, Table S2 (SI Appendix) presents an overview not of the detailed circumstances under which cocoa was produced in the plots or farms analyzed in each study, but of the range of conditions reported in the studies.

2.3 Data processing

The selected studies contained 144 pair comparisons of cocoa agroforestry systems with cocoa monocultures. We extracted all qualitative data from tables and texts, manually, and from graphs, using the software Graph Grabber 2.0, Quintessa Limited. Some articles compared one cocoa monoculture with two or more cocoa agroforestry systems, indicating that data pairs were not independent. To avoid overestimation by artificial repetition of the monoculture, we combined the data of the cocoa agroforestry systems [29] and compared the cocoa

monoculture with the mean across cocoa agroforestry systems as one data pair [30, 31]. By doing so, we reduced the number of data pairs from 144 to 93. We grouped the data into eight main categories: 1) yield (cocoa yield and total system yield); 2) economic performance (costs, revenue, net present value); 3) soil chemical properties (total soil carbon, nitrogen, phosphorous, potassium, soil organic carbon); 4) soil physical properties (bulk density, volumetric water content, mean weight diameter); 5) pests and diseases; 6) biomass of cocoa and shade trees (basal area and carbon stocks of the cocoa trees and the production system); 7) microclimate (mean, maximum and minimum temperature, relative humidity, vapor pressure deficit of the air); and 8) biodiversity (wildlife animal species and herbaceous plant species).

We calculated cocoa yield as kilograms of dry beans per hectare and year. Where only fresh weight data were provided, we converted them into dry bean weight by applying a dry bean factor of 0.35 [5]. We converted soil organic matter into soil organic carbon by dividing the former by the conversion factor 1.72 [31]. We determined total carbon stocks as the sum of the aboveground and belowground carbon of cocoa trees, shade trees, and the system (sum of cocoa trees and shade trees). When the biomass was given instead of the carbon content, we converted it into carbon stocks by multiplying the biomass by the conversion factor 0.5 [32]. When only aboveground biomass (AGB) or belowground biomass were provided, we calculated the counterpart from the given data by assuming that AGB corresponds to 87% of total biomass [33]. When only the stem diameter or basal area were given, we calculated the AGB using allometric equations for cocoa trees [34]: $\log_{10}(\text{AGB}) = (-1.625 + 2.63 * \log_{10}(\text{diameter}))$, and shade trees [35]: $\log_{10}(\text{AGB}) = (-0.834 + 2.223 * (\log_{10}(\text{diameter})))$. We calculated all economic data in USD per hectare and year. Other currencies were converted to USD using a mean exchange rate for the specific year of data collection given in the article.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

178 *2.4 Statistical analyses*

179 We used Hedge’s *g* as the effect size of our meta-analysis, which is based on the raw mean
180 difference between the mean of the cocoa agroforestry system and the mean of the cocoa
181 monoculture (the grand mean from the random effects [RE] model), standardized by the pooled
182 standard deviation (SD) across both production systems, and the sample size (*n*) of the single
183 data pair comparisons. When the SD was not provided and could not be calculated from the
184 data (i.e., from the standard error and the number of repetitions), we reassigned it as 1/10 of
185 the mean [31]. We conducted all analyses with the metafor package [36] of the R programming
186 environment, version 3.5.3 [37]. For the map (SI Appendix, Figure S4), we used the mapproj
187 package [38], and for the graphs, the ggplot2 package [39].

188
189
190 **3. Results and Discussion**

191
192 *3.1 Can cocoa agroforestry systems increase productivity?*

193 Cocoa yield in agroforestry systems is on average 75% of the cocoa production in monocultures
194 (Figure 1, Table 1). Several studies reported a negative effect of increasing shade levels on cocoa
195 yield [7, 19, 40, 41]. Cocoa tree development in agroforestry systems can be slower compared to
196 monocultures [5], which could be one of the main reasons for the often rather negative
197 perception of farmers regarding the production of cocoa under shade [42]. However, long-term
198 studies concluded that the short-term reduction of cocoa production under agroforestry is
199 compensated by the longer productive lifetime of cocoa trees grown under shade [43, 26]. Also
200 the longevity of the cocoa leaves is reduced under high solar radiation [44], which may indicate
201 a negative effect of direct solar radiation on the whole tree. The cocoa agroforestry system and
202 the cocoa monoculture compared within the data pairs had similar ages. This is important since

the effect of the tree age on cocoa yield has been reported [27]. The plantations analyzed ranged from four to 50 years, but the majority of the research was conducted in plantations up to 25 years. Data on the performance of old cocoa agroforestry systems compared to old cocoa monocultures are still scarce and often rely rather on modelling approaches than on field data [26].

Table 1. Mean values and standard deviations (SD) for a) yield; b) economic performance; c) soil chemical properties; d) soil physical properties; e) pests and diseases; f) microclimate; g) stand structural parameters in cocoa agroforestry systems and cocoa monocultures. N indicates the number of studies; levels of significance: *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, n.s.: not significant.

			Cocoa agroforestry system		Cocoa monoculture			
Group	Variable	Unit	Mean	SD	Mean	SD	N	
Yield								
	Cocoa yield	Mg ha ⁻¹	0.6	± 0.4	0.9	± 0.7	36	***
	System yield	Mg ha ⁻¹	9.8	± 9.2	0.6	± 0.4	8	*
Economic performance								
	Costs	USD ha ⁻¹ a ⁻¹	571.5	± 322.8	652.9	± 464.4	7	n.s.
	System revenue	USD ha ⁻¹ a ⁻¹	1094.3	± 594.7	1299.7	± 905.9	8	n.s.
	Net present value	USD ha ⁻¹	998.8	± 736.8	1108.9	± 729.7	4	n.s.
Soil chemical properties								
	Soil C	%	14.5	± 2.4	13.8	± 2.3	20	n.s.
	Soil N	%	1.8	± 0.7	1.7	± 0.6	22	n.s.
	Soil available P	mg kg ⁻¹	13.7	± 14.2	17.2	± 16.9	9	n.s.
	Soil available K	g kg ⁻¹	0.1	± 0.1	0.1	± 0.1	10	n.s.
	Soil organic carbon	%	1.7	± 0.5	1.7	± 0.5	8	n.s.
	pH		6.3	± 0.4	6.4	± 0.5	6	*
Soil physical properties								

1						
2						
3		Mean weight diameter	mm	1.0 ± 0.4	0.9 ± 0.2	10 n.s.
4						
5		Bulk density	g cm ³	1.3 ± 0.3	1.4 ± 0.2	4 *
6						
7		Volumetric water content	%	20.1 ± 5.4	21.8 ± 5.7	6 **
8						
9						
10	Fungal diseases					
11						
12		Frosty pod rot	%	28.8 ± 24.5	21.2 ± 16	4 n.s.
13						
14		Black pod	%	3.4 ± 2.2	3.0 ± 2.0	5 *
15						
16		Witches' broom	%	1.9 ± 1.4	3.7 ± 2.4	5 *
17						
18	Microclimate					
19						
20		Maximum temperature	°C	32.4 ± 2.5	34.7 ± 3.3	8 *
21						
22		Mean temperature	°C	18.6 ± 3.1	17.9 ± 3.4	8 ***
23						
24		Minimum temperature	°C	24.7 ± 1.8	25.0 ± 1.8	8 *
25						
26		Mean relative humidity	%	81.5 ± 16.5	80.5 ± 15.6	3 n.s.
27						
28		Vapor pressure deficit	kPa	1.1 ± 0.7	1.3 ± 0.8	4 n.s.
29						
30	Stand structural parameters					
31						
32		Basal area cocoa trees	m ² ha ⁻¹	7.7 ± 2.9	9.4 ± 3.2	22 ***
33						
34		Basal area shade trees	m ² ha ⁻¹	10.2 ± 2.2	0.2 ± 0.4	4 ***
35						
36		Total C in cocoa trees	Mg ha ⁻¹	9.5 ± 6.3	13.2 ± 6.9	30 ***
37						
38		Total C in shade trees	Mg ha ⁻¹	24.7 ± 26.3	1.0 ± 4.6	27 ***
39						
40		Total C in system	Mg ha ⁻¹	37.0 ± 28.9	14.2 ± 9.0	30 ***
41	<hr/>					

213

214 Considering all crops harvested, production in cocoa agroforestry systems amounts to

215 $9.8 \pm 9.2 \text{ Mg ha}^{-1} \text{ a}^{-1}$, which is about 10 times higher than production in cocoa monocultures

216 (Figure 1, Table 1). The lower system yield obtained in cocoa monocultures compared to cocoa

217 yield in monocultures ($0.6 \text{ vs } 0.9 \text{ Mg ha}^{-1}$) is related to different studies included as well as the

218 number of pair comparisons, i.e., we calculated the mean yield of only eight data pairs for

219 system yield, while cocoa yield is calculated from 36 data pairs. Kuyah et al. [45] obtained similar

results for other types of agroforestry systems, pointing to their potential to produce food, which improves food security of farming families [5, 25]. Production diversity also reduces dependency on one single crop and, consequently, fluctuations in prices and demand [46]. The high variability of system yields reported might be explained by the wide range of crops that can be grown in cocoa agroforestry systems (SI Appendix, Table S1). Besides the production, the commercialization of fruits can be difficult due to reduced market access or local acceptance. Banana and plantain are common products in cocoa agroforestry systems [47, 5], since they are also used for protecting the cocoa trees from direct light during the establishment phase. They produce in high quantities and may be distributed locally since they are staple crops in many regions and commonly consumed. Tuber crops like turmeric and ginger need specialized markets or export options, and uncommon fruits like araza (*Eugenia stipitata*) cannot be transported or stored easily to reach their market [48].




Figure 1. Mean difference of cocoa and total system yields in agroforestry systems compared with monocultures. The number of studies is shown in brackets; the horizontal bar shows the 95% confidence intervals; negative values indicate a higher mean value in monocultures; positive values indicate a higher mean value in cocoa agroforestry systems; levels of significance: *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, n.s.: not significant.

3.2 Can cocoa agroforestry systems sustain farmers' incomes?

Even though the total system production is higher in cocoa agroforestry systems, this is not reflected in higher revenues or net present values, neither is it in the costs (Figure 2). Cocoa, a

1
2
3 242 commodity produced mainly for export, normally reaches higher prices than its by-crops, which
4
5 243 are mainly sold in local markets, or consumed on-farm and do not contribute to farmer's income
6
7 244 [49]. The economic performance of the production systems depends on the level of
8
9
10 245 management of the plantation and on labor costs, with cocoa agroforestry systems tending to
11
12 246 have higher labor demands [24]. In the case of cocoa agroforestry systems, it also depends on
13
14 247 the planting design and selection of shade tree species [47]. Timber trees increase the net
15
16 248 present value of cocoa agroforestry systems [47]. However, their value is not always considered
17
18 249 a future benefit for farmers, due to insecure land and tree tenure and the risk of fires [42]. The
19
20
21 250 wide range of management intensities in the production systems in the different studies is likely
22
23 251 responsible for the variation in the mean differences of the economic variables analyzed
24
25 252 between cocoa agroforestry systems and monocultures (SI Appendix, Figure S1). The total costs
26
27 253 in organically managed cocoa production systems included the certification costs [24], while
28
29 254 another study included the costs for renting the land [26]. The net present value depends not
30
31 255 only on the planting design, but also on the plantation age and the timeframe that was used for
32
33 256 the calculation, e.g. a 20-year production cycle [50], or a 12-year net income flow [47].
34
35
36
37 257 The lack of a clear effect of the type of production system on variables of the economic
38
39 258 performance indicates that, despite their lower cocoa production, agroforestry systems can be
40
41
42 259 economically as viable as monocultures.



260


Figure 2. Mean difference in the economic performance of cocoa agroforestry systems compared with monocultures. The number of studies is shown in brackets; the horizontal bars show the 95% confidence intervals; negative values indicate a higher mean value in monocultures; positive values indicate a higher mean value in cocoa agroforestry systems; levels of significance: *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, n.s.: not significant.

3.3 Do cocoa agroforestry systems improve soil chemical and physical properties for cocoa production?

We found no significant differences in soil chemical properties between production systems, with the exception of higher soil pH values in monocultures (Figure 3a, Table 1). Thus, our results do not indicate a positive impact of cocoa agroforestry on soil fertility. However, we observed substantial differences between studies. For instance, studies sampling around single shade trees found positive effects on soil chemical properties [51-53]. Positive effects on soil chemical parameters could likely be achieved if the biomass resulting from pruning the shade trees were distributed across the plantation [54]. Still, shade trees are not usually pruned and distributing the biomass can be highly labor-intensive. Tree selection can ultimately influence soil chemical properties [51, 53]. Better knowledge on shade tree properties is thus important for an optimal tree selection. Finally, contrasting results between studies may also be related to different soil type, plantation age and land-use history [55], which might affect the influence of agroforestry trees on the soil [56], e.g. Mohammed et al. [57] described a slight increase in SOC with the age of the cocoa agroforestry system and a decline in monocultures.

Soils in cocoa agroforestry systems have a significantly lower volumetric water content than in monocultures (Figure 3b, Table 1). This may result in competition for water between cocoa and shade trees. However, this potentially negative effect of shade trees will only become critical if

1
2
3 285 shade trees are planted at very high densities [7], or in regions with an annual precipitation that
4
5 286 is close to the limit for cocoa production [33, 58]. This needs to be further investigated and
6
7 287 carefully considered in relation to future scenarios of decreasing or changing water availability
8
9
10 288 patterns due to climate change [59]. Potential competition for water may be reduced by
11
12
13 289 selecting shade tree species with rooting patterns complementary to that of cocoa trees [17, 60,
14
15 290 61]. However, the information available on this aspect, though crucial for providing
16
17
18 291 recommendations on tree species selection and planting patterns, is still very limited.




292
293 **Figure 3.** Mean difference in the a) soil chemical properties and b) soil physical properties of
294 cocoa agroforestry systems compared with monocultures. The number of studies is shown in
295 brackets; horizontal bars show the 95% confidence intervals; negative values indicate a higher
296 mean value in monocultures; positive values indicate a higher mean value in cocoa agroforestry
297 systems; levels of significance: *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, n.s.: not significant.

298
299 3.4 Do cocoa agroforestry systems enhance the control of pests and diseases?

1
2
3 300 Pests and diseases, and particularly fungal diseases, are a major threat to cocoa production. The
4
5 301 analysis of disease control in the different systems results in contrasting findings: frosty pod rot
6
7 302 (*Moniliophthora roreri*) is not affected by the production system, while the incidences of black
8
9 303 pod (*Phytophthora spp.*) and witches' broom (*Moniliophthora perniciosa*) are respectively higher
10
11 304 and lower in cocoa agroforestry systems than in cocoa monocultures (Figure 4, Table 1). These
12
13 305 findings are in line with previous studies showing the complex effects of shade trees on the
14
15 306 incidence of pests and diseases [7, 22, 62]. Influencing factors include: the management of the
16
17 307 system (e.g., pruning, fertilization, weeding) [63]; the specific pest and disease management
18
19 308 strategy (e.g. cultural, chemical or biological control) [62, 63]; the specific characteristics of the
20
21 309 pest or disease considered [30]; and the particular microclimatic conditions, which highly
22
23 310 depend on the structural complexity of the cocoa agroforestry system [18]. In cocoa
24
25 311 agroforestry systems, reduced light availability and wind speed, buffered temperatures and
26
27 312 increased relative humidity compared to monocultures [18] may stimulate sporulation of
28
29 313 diseases such as black pod, but also favor the presence of antagonists, e.g. for witches' broom
30
31 314 [22, 64]. The broad variance of the incidence of frosty pod rot in our analysis is not only related
32
33 315 to cocoa production systems, but also to general differences in the infestation rate. Up to 63 %
34
35 316 infected fruits were counted by Krauss and Soberanis [62] compared to a relatively low
36
37 317 infestation rate around 10 % by Armengot et al. [63] which was explained by regular elimination
38
39 318 of affected fruits before sporulation rather than application of fungicides. The single data pair
40
41 319 comparison of a broader range of pests and diseases (including cocoa swollen shoot virus
42
43 320 disease, leaf herbivory, vascular streak dieback), as well as of the total yield loss and total
44
45 321 infestation, showed contrasting results, too. This resulted in no significant difference in the
46
47 322 incidence of pests and diseases between cocoa agroforestry systems and cocoa monocultures
48
49
50
51
52
53
54 323 according to the grand mean difference over all data pairs (SI Appendix, Figure S2).

1
2
3 324 The results do not completely support the above-mentioned regulating effects of cocoa
4
5 325 agroforestry systems. However, they show that cocoa agroforestry systems compared with
6
7 326 cocoa monocultures are not prone to pests and diseases, contrary to widespread perceptions.
8
9



327
328 **Figure 4.** Mean difference in the incidence of three fungal diseases in cocoa agroforestry systems
329 and monocultures. The number of studies is shown in brackets; horizontal bars show the 95%
330 confidence intervals; negative values indicate a higher mean value in monocultures; positive
331 values indicate a higher mean value in cocoa agroforestry systems; levels of significance: *** $p <$
332 0.001, ** $p <$ 0.01, * $p <$ 0.05, n.s.: not significant.
333

334 *3.5 Do cocoa agroforestry systems support adaptation of cocoa plantations to climate change?*

335 In cocoa agroforestry systems, solar radiation is reduced by the canopy of shade trees [18],
336 which provides a more stable microclimate: the mean temperature remains lower and the daily
337 maximum and minimum temperatures are buffered compared to what happens in
338 monocultures (Figure 5, Table 1). Cocoa agroforestry systems, therefore, have the potential to
339 reduce the impact of rising mean temperatures and temperature extremes predicted for
340 producer countries [59, 65]. Consequently, they are more resilient to climate change and
341 provide more comfortable working conditions (shade and lower temperatures) than full-sun
342 monocultures [46, 66].

343 The mean relative humidity of the air and the vapor pressure deficit do not differ between
344 cocoa agroforestry systems and cocoa monocultures (Figure 5). Some studies show that the

difference in relative humidity is pronounced during the wet season, and smaller during the dry season [7]. In addition, canopy closure and stratification of cocoa agroforestry systems affect internal relative humidity [18]. Tall trees may allow more aeration below their canopy than dense tree crowns close to the cocoa canopy. Tree species selection and planting density are therefore important for shading intensity and microclimatic conditions. The actual radiation reaching cocoa agroforestry systems—which depends on location-specific factors like latitude, altitude, cloudiness, slope exposure and surrounding shading forests—needs to be considered for the selection of the shade trees [67]. It is important to highlight that microclimatic variables can be influenced and adapted to seasonal needs of cocoa tree plantations by managing shade intensity through regular shade tree pruning [18].




Figure 5. Mean difference in microclimatic variables between cocoa agroforestry systems and monocultures. The number of studies is shown in brackets; horizontal bars show the 95% confidence intervals; negative values indicate a higher mean value in monocultures; positive values indicate a higher mean value in cocoa agroforestry systems; levels of significance: *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, n.s.: not significant.

3.6 Do cocoa agroforestry systems contribute to climate change mitigation?

Cocoa trees tend to store more carbon when growing in monocultures, since they are bigger and often planted at higher densities than in agroforestry systems, as reflected in the higher basal area (Figure 6, Table 1). Shade trees are often planted at low densities, but, since they usually have larger stem diameters and grow high above the cocoa trees, they account for most of the basal area [18] and for 66% of the carbon stored in the system (Figure 6, Table 1). The total carbon (aboveground and root biomass) stored in an agroforestry system, including both the cocoa and the shade trees, is, on average, 2.5 times higher than in a monoculture (Figure 6, Table 1). Thus, cocoa agroforestry systems have greater potential for climate change mitigation than cocoa monocultures due to a higher carbon sequestration potential. Carbon payments are often mentioned as a potential incentive for farmers to plant trees. However, widely accessible systems directing these payments to cocoa producers are rarely found, and the payments tend to be too low to incentivize the planting of shade trees [68].





Figure 6. Mean difference in stand structural parameters between cocoa agroforestry systems and monocultures. The number of studies is shown in brackets; horizontal bars show the 95% confidence intervals; negative values indicate a higher mean value in monocultures; positive values indicate a higher mean value in cocoa agroforestry systems; levels of significance: *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, n.s.: not significant; Total C: carbon stock of aboveground and root biomass.

3.7 Do cocoa agroforestry systems contribute to biodiversity conservation?

Only five articles directly compare the species richness of five animal groups [7, 46, 69, 70, 71] in cocoa agroforestry systems and monocultures. Therefore, we prefer to show the results for each taxon and study rather than only the mean difference. The mean difference in the grand mean over all animal taxa (RE Model) between cocoa agroforestry systems and cocoa monocultures shows a significantly higher number of species in agroforestry systems (Figure 7a). This positive effect of shade trees on biodiversity is consistent with other studies that have investigated the value of cocoa agroforests for the conservation of biodiversity [19, 72-74]. The effect of shade on animal biodiversity depends amongst others on the taxa under consideration [73], but also on the management, diversity, and complexity of the particular cocoa agroforestry system [75]. Agroforestry systems have also been reported to provide habitat for functionally more diverse species communities because they are structurally more complex and diverse than monocultures [5, 20, 76] (SI Appendix, Table S1).

The scarcity of available data on the diversity of herbaceous species in cocoa agroforestry systems and cocoa monocultures (two data pair analyses from only one publication [21]) prevents us from generalizing the results by calculating the grand mean difference (RE Model) (Figure 7b). However, the data imply that the effect on herbaceous species richness might depend heavily on farm management practices, e.g., use of agrochemicals or cover crops, availability of light, and weeding interventions [20, 21].






Figure 7. Comparison of a) animal and b) herbaceous species richness in cocoa agroforestry systems and monocultures shown by single studies and the mean difference between studies (RE Model). Horizontal bars show the 95% confidence intervals; negative values indicate a higher mean value in monocultures; positive values indicate a higher mean value in cocoa agroforestry systems.

4. Conclusions

This meta-analysis indicates that cocoa agroforestry systems have the potential to compete with cocoa monocultures in terms of economic performance, and to outperform them in crucial system services such as adaptation to climate change and carbon sequestration, as well as in total system yields. With only five articles on biodiversity conservation in cocoa agroforestry systems and cocoa monocultures, and six on the incidence of pests and diseases, we identified a knowledge gap for these two topics in cocoa production. Although this prevents us from generalizing the results, it is possible to infer that cocoa agroforestry systems tend to have a similar or even better performance than monocultures in most of the evaluated parameters.

Our results underline the need for promoting cocoa agroforestry systems to improve the sustainability of the cocoa sector. Despite all the above-mentioned benefits of cocoa agroforestry systems, the lower cocoa yield might still be one of the most relevant factors hindering a broader adoption of diversified production systems. In this sense, further research focused on increasing cocoa yields in agroforestry systems is crucial, e.g., breeding for shade tolerant varieties or adapted management practices to increase pollination rates. However, for

promoting cocoa agroforestry systems, building and enabling access to new alternative markets and value chains for agroforestry products is also crucial, as is compensating farmers for cocoa yield reductions through fair prices for sustainable cocoa production or carbon storage. The promotion and support of cocoa agroforestry systems by the cocoa industry can therefore contribute to meeting its sustainability goals.

The high heterogeneity and wide range of climatic and edaphic conditions, management strategies, planting densities and shade tree species encountered in the literature included in this meta-analysis suggest that a global recommendation for shade levels or shade tree species would not be accurate. Rather, local and context-specific recommendations for cocoa agroforestry design and management considering socio-economic factors should be developed for the implementation of sustainable and feasible cocoa production systems. Knowledge gaps regarding, for instance, detailed species-specific information on shade trees need to be addressed. The role of different shade trees on soil nutrient dynamics, including competition and synergies for resources, needs to be elucidated. This needs to be context specific, considering different soil types and land-use histories. Finally, management strategies, pricing policies, cultural and social services, and livelihood aspects deserve further attention in future research.

Acknowledgments

We wish to thank the authors of the studies that are the basis of this meta-analysis. The study was funded by the Swiss Platform for Sustainable Cocoa. Laura Armengot was supported by the Syscom programme, financed by the Swiss Agency for Development and Cooperation (SDC), the

Liechtenstein Development Service (LED), the Biovision Foundation for Ecological Development, and the Coop Sustainability Fund.

Data availability statement

No new data were created or analyzed in this study.

References

[1] FAOSTAT, FAOSTAT. 2020, Food and Agricultural Organization of the United Nations

[2] Hütz-Adams F, Huber C, Knoke I, Morazan P, and Mürlebach M, Strengthening the competitiveness of cocoa production and improving the income of cocoa producers in West and Central Africa, S.e. V., Editor. 2016: Bonn

[3] Shabandeh M, Global cocoa processing market value 2017-2023. 2018: Statista

[4] Almeida A and Valle R 2007 Ecophysiology of the cacao tree *Brazilian Journal of Plant Physiology* **19** 425-448

[5] Schneider M, Andres C, Trujillo G, Alcon F, Amurrio P, Perez E, Weibel F, and Milz J 2016 Cocoa and total system yields of organic and conventional agroforestry vs. monoculture systems in a long-term field trial in Bolivia *Experimental Agriculture* **53** 351-374

[6] Ruf F and Schroth G 2004 *Chocolate forests and monocultures: a historical review of cocoa growing and its conflicting role in tropical deforestation and forest conservation*, in *Agroforestry and biodiversity conservation in tropical landscapes* G. Schroth, et al., Editors. (Washington, DC: Island Press)

[7] Blaser WJ, Oppong J, Hart SP, Landolt J, Yeboah E, and Six J 2018 Climate-smart sustainable agriculture in low-to-intermediate shade agroforests *Nature Sustainability* **1** 234-239

[8] Somarriba E, Cerda R, Orozco L, Cifuentes M, Dávila H, Espin T, Mavisoy H, Ávila G, Alvarado E, Poveda V, Astorga C, Say E, and Deheuvels O 2013 Carbon stocks and cocoa yields in agroforestry systems of Central America *Agriculture, Ecosystems & Environment* **173** 46-57

[9] Takyi SA, Amponsah O, Inkoom DKB, and Azunre GA 2019 Sustaining Ghana's cocoa sector through environmentally smart agricultural practices: an assessment of the environmental impacts of cocoa production in Ghana *Africa Review* **11** 172-189

[10] Pérez-Neira D, Schneider M, and Armengot L 2020 Crop-diversification and organic management increase the energy efficiency of cacao plantations *Agricultural Systems* **177** 102711

[11] Tothmihaly A 2017 *How low is the price elasticity in the global cocoa market?* Global Food Discussion Papers. Vol. 102. (Göttingen: Georg-August-Universität Göttingen, Research Training Group (RTG))

- 492 [12] Jumiya S, Arsyad M, Rajindra, Pulubuhu DAT, and Hadid A 2018 Cocoa based
 493 agroforestry: An economic perspective in resource scarcity conflict era *IOP Conference*
 494 *Series: Earth and Environmental Science* **157** 012009
- 495 [13] Beillouin D, Ben-Ari T, and Makowski D 2019 Evidence map of crop diversification
 496 strategies at the global scale *Environmental Research Letters* **14** 123001
- 497 [14] Jacobi J, Rist S, and Altieri MA 2017 Incentives and disincentives for diversified
 498 agroforestry systems from different actors' perspectives in Bolivia *International Journal*
 499 *of Agricultural Sustainability* **15** 365-379
- 500 [15] Nestlé Nestlé cocoa plan 2016 15 February 2020 <https://www.nestlecocoa.com/>
- 501 [16] Lindt & Sprüngli Sustainably Sourced 2008 15 February 2020 [https://www.lindt-](https://www.lindt-spruengli.com/sustainability/)
 502 [spruengli.com/sustainability/](https://www.lindt-spruengli.com/sustainability/)
- 503 [17] Tscharntke T, Clough Y, Bhagwat SA, Buchori D, Faust H, Hertel D, Hölscher D, Juhrbandt
 504 J, Kessler M, and Perfecto I 2011 Multifunctional shade-tree management in tropical
 505 agroforestry landscapes—a review *Journal of Applied Ecology* **48** 619-629
- 506 [18] Niether W, Armengot L, Andres C, Schneider M, and Gerold G 2018 Shade trees and tree
 507 pruning alter throughfall and microclimate in cocoa (*Theobroma cacao* L.) production
 508 systems *Annals of Forest Science* **75** 38
- 509 [19] Clough Y, Putra DD, Pitopang R, and Tscharntke T 2009 Local and landscape factors
 510 determine functional bird diversity in Indonesian cacao agroforestry *Biological*
 511 *Conservation* **142** 1032-1041
- 512 [20] Clough Y, Barkmann J, Juhrbandt J, Kessler M, Wanger TC, Anshary A, Buchori D, Ciczuzza
 513 D, Darras K, and Putra DD 2011 Combining high biodiversity with high yields in tropical
 514 agroforests *Proceedings of the National Academy of Sciences* **108** 8311-8316
- 515 [21] Marconi L and Armengot L 2020 Complex agroforestry systems against biotic
 516 homogenization: The case of plants in the herbaceous stratum of cocoa production
 517 systems *Agriculture, Ecosystems & Environment* **287** 106664
- 518 [22] Mortimer R, Saj S, and David C 2018 Supporting and regulating ecosystem services in
 519 cacao agroforestry systems *Agroforestry Systems* **92** 1639-1657
- 520 [23] Altieri MA 1999 *The ecological role of biodiversity in agroecosystems*, in *Invertebrate*
 521 *biodiversity as bioindicators of sustainable landscapes*: Elsevier). p. 19-31
- 522 [24] Armengot L, Barbieri P, Andres C, Milz J, and Schneider M 2016 Cacao agroforestry
 523 systems have higher return on labor compared to full-sun monocultures *Agronomy for*
 524 *sustainable development* **36** 70
- 525 [25] Jacobi J 2016 Agroforestry in Bolivia: opportunities and challenges in the context of food
 526 security and food sovereignty *Environmental Conservation* **43** 307-316
- 527 [26] Obiri BD, Bright GA, McDonald MA, Anglaaere LC, and Cobbina J 2007 Financial analysis
 528 of shaded cocoa in Ghana *Agroforestry systems* **71** 139-149
- 529 [27] Wessel M, and Quist-Wessel PMF 2015 Cocoa production in West Africa, a review and
 530 analysis of recent developments *NJAS - Wageningen Journal of Life Sciences* **74-75** 1-7
- 531 [28] De Beenhouwer M, Aerts R, and Honnay O 2013 A global meta-analysis of the
 532 biodiversity and ecosystem service benefits of coffee and cacao agroforestry
 533 *Agriculture, Ecosystems & Environment* **175** 1-7
- 534 [29] Higgins JP and Green S, *Cochrane handbook for systematic reviews of interventions*.
 535 Chichester, England; Hoboken. 2008, NJ: Wiley-Blackwell
- 536 [30] Andres C, Blaser WJ, Dzahini-Obiatey HK, Ameyaw GA, Domfeh OK, Awiagah MA,
 537 Gattinger A, Schneider M, Offei SK, and Six J 2018 Agroforestry systems can mitigate the
 538 severity of cocoa swollen shoot virus disease *Agriculture, Ecosystems & Environment*
 539 **252** 83-92

- [31] Gattinger A, Muller A, Haeni M, Skinner C, Fliessbach A, Buchmann N, Mäder P, Stolze M, Smith P, and Scialabba NE-H 2012 Enhanced top soil carbon stocks under organic farming *Proceedings of the National Academy of Sciences* **109** 18226-18231
- [32] Pearson T, Harris N, Shoch D, and Brown S 2005 A sourcebook of methods and procedures for monitoring and reporting anthropogenic greenhouse gas emissions and removals associated with deforestation, gains and losses of carbon stocks in forests remaining forests, and forestation *Global Observation of Forest and Land Cover Dynamics*
- [33] Zuidema PA, Leffelaar PA, Gerritsma W, Mommer L, and Anten NP 2005 A physiological production model for cocoa (*Theobroma cacao*): model presentation, validation and application *Agricultural Systems* **84** 195-225
- [34] Andrade H, Segura MA, Somarriba Chávez E, and Villalobos Rodríguez M 2008 Valoración biofísica y financiera de la fijación de carbono por uso del suelo en fincas cacaoteras indígenas de Talamanca, Costa Rica *Agroforestería en las Américas* **46**
- [35] Segura M and Andrade Castañeda HJ 2008 ¿Cómo construir modelos alométricos de volumen, biomasa o carbono de especies leñosas perennes?
- [36] Viechtbauer W 2010 Conducting meta-analyses in R with the metafor package *Journal of statistical software* **36** 1-48
- [37] R Core Team, R: A language and environment for statistical computing, R.F.f.S. Computing, Editor. 2019: Vienna
- [38] McIlroy D, Mapproj: Map Projections in *Package for R by Ray Brownrigg, Thomas P Minka and transition to Plan 9 codebase by Roger Bivand*. 2018
- [39] Wickham H 2016 *ggplot2: elegant graphics for data analysis*. Springer
- [40] Pérez Neira D 2016 Energy efficiency of cacao agroforestry under traditional and organic management *Agronomy for Sustainable Development* **36** 49
- [41] Riedel J, Kägi N, Armengot L, and Schneider M 2019 Effects of rehabilitation pruning and agroforestry on cacao tree development and yield in an older full-sun plantation *Experimental Agriculture* **55** 849-865
- [42] Ruf FO 2011 The myth of complex cocoa agroforests: the case of Ghana *Human ecology* **39** 373-388
- [43] Ahenkorah Y, Akrofi G, and Adri A 1974 The end of the first cocoa shade and manurial experiment at the Cocoa Research Institute of Ghana *Journal of Horticultural Science* **49** 43-51
- [44] Miyaji K-I, Da Silva WS, and Alvim PdT 1997 Longevity of leaves of a tropical tree, *Theobroma cacao*, grown under shading, in relation to position within the canopy and time of emergence *The New Phytologist* **135** 445-454
- [45] Kuyah S, Whitney CW, Jonsson M, Sileshi GW, Öborn I, Muthuri CW, and Luedeling E 2019 Agroforestry delivers a win-win solution for ecosystem services in sub-Saharan Africa. A meta-analysis *Agronomy for sustainable development* **39**
- [46] Jacobi J, Schneider M, Bottazzi P, Pillco M, Calizaya P, and Rist S 2015 Agroecosystem resilience and farmers' perceptions of climate change impacts on cocoa farms in Alto Beni, Bolivia *Renewable Agriculture and Food Systems* **30** 170-183
- [47] Ramirez O, Somarriba E, Ludewigs T, and Ferreira P 2001 Financial returns, stability and risk of cacao-plantain-timber agroforestry systems in Central America *Agroforestry Systems* **51** 141-154
- [48] Niether W, Schneidewind U, Fuchs M, Schneider M, and Armengot L 2019 Below-and aboveground production in cocoa monocultures and agroforestry systems *Science of the Total Environment* **657** 558-567

- [49] Cerda R, Deheuvels O, Calvache D, Niehaus L, Saenz Y, Kent J, Vilchez S, Villota A, Martinez C, and Somarriba E 2014 Contribution of cocoa agroforestry systems to family income and domestic consumption: looking toward intensification *Agroforestry systems* **88** 957-981
- [50] Gockowski J, Afari-Sefa V, Sarpong DB, Osei-Asare YB, and Agyeman NF 2013 Improving the productivity and income of Ghanaian cocoa farmers while maintaining environmental services: what role for certification? *International Journal of Agricultural Sustainability* **11** 331-346
- [51] Isaac ME, Timmer VR, and Quashie-Sam S 2007 Shade tree effects in an 8-year-old cocoa agroforestry system: biomass and nutrient diagnosis of *Theobroma cacao* by vector analysis *Nutrient cycling in agroecosystems* **78** 155-165
- [52] Blaser WJ, Oppong J, Yeboah E, and Six J 2017 Shade trees have limited benefits for soil fertility in cocoa agroforests *Agriculture, Ecosystems & Environment* **243** 83-91
- [53] Wartenberg AC, Blaser WJ, Roshetko JM, Van Noordwijk M, and Six J 2019 Soil fertility and *Theobroma cacao* growth and productivity under commonly intercropped shade-tree species in Sulawesi, Indonesia *Plant and Soil* 1-18
- [54] Schneidewind U, Niether W, Armengot L, Schneider M, Sauer D, Heitkamp F, and Gerold G 2019 Carbon stocks, litterfall and pruning residues in monoculture and agroforestry cacao production systems *Experimental Agriculture* **55** 452-470
- [55] Martin DA, Osen K, Grass I, Hölscher D, Tschantke T, Wurz A, and Kreft H 2020 Land-use history determines ecosystem services and conservation value in tropical agroforestry *Policy Perspective*, online first
- [56] Centenaro G, Hudek C, Zanella A, and Crivellaro A 2018 Root-soil physical and biotic interactions with a focus on tree root systems: A review *Applied soil ecology* **123** 318-327
- [57] Mohammed AM, Robinson JS, Midmore D, and Verhoef A 2016 Carbon storage in Ghanaian cocoa ecosystems *Carbon Balance and Management* **11** 1-8
- [58] Abdulai I, Vaast P, Hoffmann MP, Asare R, Jassogne L, Van Asten P, Rötter RP, and Graefe S 2018 Cocoa agroforestry is less resilient to sub-optimal and extreme climate than cocoa in full sun *Global Change Biology* **24** 273-286
- [59] Läderach P, Martinez-Valle A, Schroth G, and Castro N 2013 Predicting the future climatic suitability for cocoa farming of the world's leading producer countries, Ghana and Côte d'Ivoire *Climatic change* **119** 841-854
- [60] Jose S 2009 Agroforestry for ecosystem services and environmental benefits: an overview *Agroforestry systems* **76** 1-10
- [61] Niether W, Schneidewind U, Armengot L, Adamtey N, Schneider M, and Gerold G 2017 Spatial-temporal soil moisture dynamics under different cocoa production systems *Catena* **158** 340-349
- [62] Krauss U and Soberanis W 2001 Rehabilitation of diseased cacao fields in Peru through shade regulation and timing of biocontrol measures *Agroforestry Systems* **53** 179-184
- [63] Armengot L, Ferrari L, Milz J, Velásquez F, Hohmann P, and Schneider M 2020 Cacao agroforestry systems do not increase pest and disease incidence compared with monocultures under good cultural management practices *Crop Protection* **130** 105047
- [64] Evans H, Krauss U, Rios R, Zeceovich A, and Arevalo-Gardini E 1998 Cocoa in Peru *Cocoa Growers' Bulletin* 7-22
- [65] Schroth G, Läderach P, Martinez-Valle AI, Bunn C, and Jassogne L 2016 Vulnerability to climate change of cocoa in West Africa: Patterns, opportunities and limits to adaptation *Science of the Total Environment* **556** 231-241

- [66] Jacobi J, Andres C, Schneider M, Pillco M, Calizaya P, and Rist S 2014 Carbon stocks, tree diversity, and the role of organic certification in different cocoa production systems in Alto Beni, Bolivia *Agroforestry systems* **88** 1117-1132
- [67] Suarez-Salazar JC, Melgarejo LM, Casanoves F, Di Rienzo JA, DaMatta FM, and Armas C 2018 Photosynthesis limitations in cacao leaves under different agroforestry systems in the Colombian Amazon *PloS one* **13**
- [68] Middelndorp RS, Vanacker V, and Lambin EF 2018 Impacts of shaded agroforestry management on carbon sequestration, biodiversity and farmers income in cocoa production landscapes *Landscape Ecology* **33** 1953-1974
- [69] Felicitas AC, Hervé BD, Ekesi S, Akutse KS, Djuidéu CT, Meupia MJ, and Babalola OO 2018 Consequences of shade management on the taxonomic patterns and functional diversity of termites (Blattodea: Termitidae) in cocoa agroforestry systems *Ecology and Evolution* **8** 11582-11595
- [70] Young AM 1983 Seasonal Differences in Abundance and Distribution of Cocoa-Pollinating Midges in Relation to Flowering and Fruit Set Between Shaded and Sunny Habitats of the La Lola Cocoa Farm in Costa Rica *Journal of Applied Ecology* **20** 801-831
- [71] Bigger M 1981 Observations on the insect fauna of shaded and unshaded Amelonado cocoa Bulletin of Entomological Research **71** 107-119
- [72] Abrahamczyk S, Kessler M, Putra DD, Waltert M, and Tscharntke T 2008 The value of differently managed cacao plantations for forest bird conservation in Sulawesi, Indonesia *Bird Conservation International* **18** 349-362
- [73] Bisseleua D, Missoup A, and Vidal S 2009 Biodiversity conservation, ecosystem functioning, and economic incentives under cocoa agroforestry intensification *Conservation biology* **23** 1176-1184
- [74] Schulze CH, Waltert M, Kessler PJ, Pitopang R, Veddeler D, Mühlenberg M, Gradstein SR, Leuschner C, Steffan-Dewenter I, and Tscharntke T 2004 Biodiversity indicator groups of tropical land-use systems: comparing plants, birds, and insects *Ecological applications* **14** 1321-1333
- [75] Perfecto I and Vandermeer J 2008 Biodiversity conservation in tropical agroecosystems
- [76] Tscharntke T, Sekercioglu CH, Dietsch TV, Sodhi NS, Hoehn P, and Tylianakis JM 2008 Landscape constraints on functional diversity of birds and insects in tropical agroecosystems *Ecology* **89** 944-951